

A framework for incorporating MJO and ENSO information into CPC probabilistic extended range forecasts

Nat Johnson¹

Emily Riddle^{2,3}, Marshall Stoner², Michelle L'Heureux², Dan Collins², Steven Feldstein⁴, and Shang-Ping Xie¹

¹*International Pacific Research Center, University of Hawai'i*

²*NOAA/NCEP Climate Prediction Center*

³*Wyle Information Systems*

⁴*Pennsylvania State University*

Accomplishments over the past year:

- Diagnosed the influence of the MJO and ENSO on the dominant wintertime circulation patterns over the extended Pacific/North America region
- Assessed the ability of the CFSv2 to capture these relationships (Emily Riddle's talk)

Objectives for the coming year:


- Develop other indices that capture relationships among tropical convection, storm track variability and midlatitude weather predictability
- Test a framework for incorporating MJO and ENSO information into extended range forecasts

K-means cluster analysis of 500 hPa geopotential height


Dataset: 500 hPa heights from NCEP/NCAR Reanalysis

Time Range: November – March (Winter months) 1979-2011


Averaging: 7-Day running mean




1 15%




2 19%




3 10%




4 12%




5 12%



6 14%



7 19%



Mean 500 hPa Geopotential height anomalies (m)



The MJO strongly impacts three cluster patterns.

Cluster 4

Cluster 6



Cluster 7

MJO enhanced convection (blue)



ENSO strongly modulates the MJO impact on the cluster frequencies.


Cluster 6



- Cluster 6 strongly enhanced after phases 5 and 6 of the MJO during El Niño but suppressed for most MJO phases during La Niña.
- Statistically significant anomalous frequencies for lags of 10-30 days


But how can CPC incorporate this sort of information into their extended range forecast products?

CPC 8-14 day (Oct 6-12) Outlook: 500 hPa heights




THE OFFICIAL 8-14 DAY HEIGHT [PROG](#) CONSISTS OF: **5 PERCENT** OF TODAY'S OPERATIONAL 0Z [GFS](#) CENTERED ON DAY 11...**5 PERCENT** OF TODAY'S OPERATIONAL 6Z [GFS](#) CENTERED ON DAY 11...**20 PERCENT** OF TODAY'S 0Z [GFS ENSEMBLE](#) MEAN CENTERED ON DAY 11...**20 PERCENT** OF TODAY'S 6Z [GFS ENSEMBLE](#) MEAN CENTERED ON DAY 11...AND **50 PERCENT** OF TODAY'S 0Z EUROPEAN ENSEMBLE MEAN CENTERED ON DAY 11.

Temperature



Precipitation



FORECAST CONFIDENCE FOR THE 6-10 DAY PERIOD: NEAR AVERAGE, 3 ON A SCALE OF 1 TO 5

Merging forecast guidance through Bayes' theorem

$$P(\boldsymbol{\theta} | C_j) \propto P(\boldsymbol{\theta})P(C_j | \boldsymbol{\theta})$$

$[\theta_1, \theta_2, \dots, \theta_K]'$

Occurrence probability for
cluster patterns 1-K

Model forecast of
cluster pattern j

Prior probability of $\boldsymbol{\theta}$

- Incorporates historical
MJO and ENSO
information

- May incorporate model
consistency from
previous forecasts

Conditional

probability of C_j ,
given $\boldsymbol{\theta}$

- Based on
archived model
forecasts or
reforecasts


- *posterior probability* of $\boldsymbol{\theta}$ can be used to construct model forecast field weights in extended range forecast
- Extended range forecast explicitly combines MJO and ENSO information with model consistency and skill information

Intended benefits of a Bayesian approach

- Less subjective model weighting that combines several different sources of forecast information (better forecasts in the 6-14 day period?)
- Potential to quantify forecast uncertainty more rigorously
- Identification of “forecasts of opportunity” beyond 14-day period
- Flexible framework that can be modified easily

Wintertime 500 hPa Geopotential Height Forecast Skill over the Extended Pacific/North America region

Wintertime El Nino and MJO Phase 1



Summary

- The combined influence of the MJO and ENSO significantly impacts the frequency of the dominant wintertime circulation patterns over the extended Pacific/North America region.
- Statistically significant anomalous frequencies of geopotential height cluster patterns extend to lags of up to 40 days, which suggests promise of useful predictability information in the one- to four-week time period.
- Over the next year we plan to combine these results with model forecast performance information (starting with CFSv2) to test a Bayesian framework for extended range forecasts.